

Mammographic Density

John Hopper

Professor & Director (Research)

Centre for Epidemiology and Biostatistics

Melbourne School of Population and Global Health

The University of Melbourne

- AMDRF – ability to collect mammos and measure MD
- AMDTSS – ~3,000, DNA for >2,000
- ABCFR – ~1,000, DNA for most
- MCCR – 4,000 (800 cases), DNA for all
- TAHS MD study – 500 with growth measured from 7-15
- Cambridge/Norwich – case-control – no BMI
- Korean Case-control Studies – Samsung, Asan - ?DNA
- Hawaiian/Japanese studies
- Vietnam MD study
- **lifepool** – including ASEAN and A-J studies

How Used

- Genetic and environmental (measured and unmeasured) determinants of MD
- MD as a risk factor (MCCS, ABCFS, AMDTTSS, Korean and other studies)
- MD as a biomarker to assess (early-life) exposures
- New SMARTs – Alto Cumulus, Cirro Cumulus, Cirrus
- Inform biological studies
- Insights into variation in risks with population sub-types

- Cf. with other measures e.g. AutoDensity
- Integration with risk models, including familial risk profile measures (BOADICEA, H score)
- Refinement and implementation of Cirrus into BreastScreen St Vincents
- BreastScreen Victoria studies of screening intervals
- Improve cost-effectiveness of mammographic screening